

# Math 501: Combinatorics

## Homework 6

Recall that you must hand in a subset of the problems for which deleting any problem makes the total score less than 10. The maximum possible score on this homework is 10 points. See the syllabus for scoring details.

NOTE: All problems from Stanley are from the second edition, which is available for free on his website here: <http://www-math.mit.edu/~rstan/ec/ec1.pdf>

### Problems

1. (1) [1 point] Using the two formulas about counting parking functions we proved in class, show that

$$\sum_h \binom{n}{h_1, h_2, \dots, h_n} = (n+1)^{n-1},$$

where the sum ranges over all sequences  $h$  of positive integers such that  $h_1 + \dots + h_i \geq i$  for all  $i$ .

2. (1+) [2 points] Find a closed formula for the exponential generating function of the sequence  $a_n = n^2$ . That is, simplify

$$\sum_{n=0}^{\infty} \frac{n^2}{n!} x^n.$$

3. (1+) [2 points] Read the definition of alternating permutations in Section 1.4 of Stanley, and write out all 16 alternating permutations of size 5. Then, read Proposition 1.6 and its proof, and either (a) comment on how awesome it is and what you learned from it, OR (b) point out something you didn't fully understand about the proof.
4. (2+) [4 points] Use generating functions to prove the identity

$$\sum_{k=0}^n \binom{2k}{k} \binom{2(n-k)}{(n-k)} = 4^n.$$

(Hint: The generating function for the Catalan numbers may come in handy!)

5. (1+) [2 points] A **binary tree** of length  $n$  is a graph on  $n$  vertices constructed recursively as follows. The empty set is a binary tree of length 0. Otherwise a binary tree has a *root vertex*  $v$ , a *left subtree*  $T_1$ , and a *right subtree*  $T_2$ , each of which is also a binary tree having a root vertex. We draw the root vertex at the top and draw an edge going down and left from  $v$  and to the root vertex of  $T_1$ , and an edge going down and right from  $v$  to the root vertex of  $T_2$ , and draw each of  $T_1$  and  $T_2$  recursively in the same manner.

Prove that the number of binary trees on  $n$  vertices is the  $n$ th Catalan number  $C_n$ . (Hint: Show they satisfy the same recursion we proved in class for Dyck paths.)

6. (2-) [3 points] A **triangulation** of a convex  $n+2$ -gon is a collection of  $n-1$  diagonals that do not intersect each other. Show that the number of triangulations of a convex  $n+2$ -gon is the  $n$ th Catalan number  $C_n$ . (Hint: Again, show they satisfy the recursion.)
7. (2+) (4 points) Find an explicit bijection between the ways of fully parenthesizing a product of  $n+1$  factors, and the Dyck paths of length  $n$ .
8. (2+) [4 points] A **derangement** of  $[n]$  is a permutation  $\pi \in S_n$  having no fixed points, that is,  $\pi_i \neq i$  for all  $i$ . Let  $D_n$  be the number of derangements of  $[n]$ . Prove that

$$\sum_{n=0}^{\infty} \frac{D_n}{n!} x^n = \frac{e^{-x}}{1-x}.$$

9. (2) [3 points] Stanley chapter 1 problem 101.
10. (5) [ $\infty$  points] Let  $P$  be a Dyck path of height  $n$ , that is, a path from  $(0, 0)$  to  $(n, n)$  using only up and right unit steps and staying weakly above the diagonal  $x = y$ . Define the *bounce path* of  $P$  as follows: starting at  $(0, 0)$ , draw a path upwards until it reaches the beginning of a rightward step in  $P$ . Then turn right and continue drawing the path until it hits the diagonal  $y = x$ . Then turn upwards again until it hits the beginning of another rightward step, and so on. If the bounce path hits the diagonal at points  $(0, 0) = (j_0, j_0), (j_1, j_1), \dots, (j_k, j_k) = (n, n)$ , define the *bounce* of  $P$  to be

$$\text{bounce}(P) = \sum_{i=1}^{k-1} (n - j_i).$$

Also define the *area* of  $P$  to be the number of complete unit squares between  $P$  and the line  $x = y$ . Then define

$$C_n(q, t) = \sum_{D \in \mathcal{P}_n} q^{\text{area}(D)} t^{\text{bounce}(D)}$$

where  $\mathcal{P}_n$  is the set of all Dyck paths of height  $n$ .

The quantity  $C_n(q, t)$  is a natural  $q, t$ -analog of the Catalan numbers that arises in the study of certain doubly graded  $S_n$ -modules. Due to this algebraic connection, it is known that

$$C_n(q, t) = C_n(t, q).$$

Find a combinatorial proof of this symmetry relation.

11. (5) [ $\infty$  points] It is known that the “super Catalan numbers”

$$S(m, n) = \frac{(2m)!(2n)!}{m!n!(m+n)!}$$

have the following properties:

- $C_n = \frac{1}{2}S(1, n)$
- $\binom{2n}{n} = S(0, n)$
- $\frac{1}{2}S(m, n)$  is an integer unless  $(m, n) = (0, 0)$

Find a combinatorial interpretation of  $\frac{1}{2}S(m, n)$ .